

Literaturliste zum aEEG

des CSM Neuro Screens™

- (2) Rundgren M, Rosen I, Friberg H. Amplitude-integrated EEG (aEEG) predicts outcome after cardiac arrest and induced hypothermia. *Intensive Care Med* 2006 June;32(6):836-42.
- (3) Wittman JJ, Jr., Hirsch LJ. Continuous electroencephalogram monitoring in the critically ill. *Neurocrit Care* 2005;2(3):330-41.
- (4) Toet MC, van der MW, de Vries LS et al. Comparison between simultaneously recorded amplitude integrated electroencephalogram (cerebral function monitor) and standard electroencephalogram in neonates. *Pediatrics* 2002 May;109(5):772-9.
- (5) Vespa PM, Nenov V, Nuwer MR. Continuous EEG monitoring in the intensive care unit: early findings and clinical efficacy. *J Clin Neurophysiol* 1999 January;16(1):1-13.
- (6) Hellstrom-Westas L, Klette H, Thorngren-Jerneck K, Rosen I. Early prediction of outcome with aEEG in preterm infants with large intraventricular hemorrhages. *Neuropediatrics* 2001 December;32(6):319-24.
- (7) Claassen J, Mayer SA, Kowalski RG et al. Detection of electrographic seizures with continuous EEG monitoring in critically ill patients. *Neurology* 2004 May 25;62(10):1743-8.
- (8) Whitelaw A, White RD. Training neonatal staff in recording and reporting continuous electroencephalography. *Clin Perinatol* 2006 September;33(3):667-77, vii.
- (9) Young GB, Jordan KG, Doig GS. An assessment of nonconvulsive seizures in the intensive care unit using continuous EEG monitoring: an investigation of variables associated with mortality. *Neurology* 1996 July;47(1):83-9.
- (10) Towne AR, Waterhouse EJ, Boggs JG et al. Prevalence of nonconvulsive status epilepticus in comatose patients. *Neurology* 2000 January 25;54(2):340-5.
- (11) Hellstrom-Westas L RldVLaGG. Amplitude-integrated EEG Classification and Interpretation in Preterm and Term Infants. *Neoreviews* 2006;Feb.7(2):76-87. 7[2], 76-87. 2007. *NeoReviews*.
- (12) Prior PF, Maynard DE, Sheaff PC et al. Monitoring cerebral function: clinical experience with new device for continuous recording of electrical activity of brain. *Br Med J* 1971 June 26;2(5764):736-8.
- (13) Rennie JM, Chorley G, Boylan GB et al. Non-expert use of the cerebral function monitor for neonatal seizure detection. *Arch Dis Child Fetal Neonatal Ed* 2004 January;89(1):F37-F40.
- (14) Laver S, Farrow C, Turner D, Nolan J. Mode of death after admission to an intensive care unit following cardiac arrest. *Intensive Care Med* 2004 November;30(11):2126-8.

- (15) Lowenstein DH, Alldredge BK. Status epilepticus. *N Engl J Med* 1998 April 2;338(14):970-6.
- (16) Sirven JI, Waterhouse E. Management of status epilepticus. *Am Fam Physician* 2003 August 1;68(3):469-76.
- (17) Garzon E, Fernandes RM, Sakamoto AC. Analysis of clinical characteristics and risk factors for mortality in human status epilepticus. *Seizure* 2003 September;12(6):337-45.
- (18) Chapman MG, Smith M, Hirsch NP. Status epilepticus. *Anaesthesia* 2001 July;56(7):648-59.
- (19) Doyle PW, Matta BF. Burst suppression or isoelectric encephalogram for cerebral protection: evidence from metabolic suppression studies. *Br J Anaesth* 1999 October;83(4):580-4.
- (20) Leistriz L, Jager H, Schelenz C et al. New approaches for the detection and analysis of electroencephalographic burst-suppression patterns in patients under sedation. *J Clin Monit Comput* 1999 August;15(6):357-67.
- (21) DeLorenzo RJ, Waterhouse EJ, Towne AR et al. Persistent nonconvulsive status epilepticus after the control of convulsive status epilepticus. *Epilepsia* 1998 August; 39(8):833-40.
- (22) Treatment of convulsive status epilepticus. Recommendations of the Epilepsy Foundation of America's Working Group on Status Epilepticus. *JAMA* 1993 August 18;270(7):854-9.
- (23) Thornberg E, Thiringer K . Normal pattern of the cerebral function monitor trace in term and preterm neonates. *Acta Paediatr Scand.* 1990;79:20-25.
- (24) Hellström-Westas L, Rosen I, Svenningsen NW. Silent seizures in sick infants in early life. *Acta Paediatr Scand* 1985; 74: 741-748.
- (25) Hellström-Westas L, Rosen I, Svenningsen NW . Predictive value of early continuous amplitude integrated EEG recordings on outcome after severe birth asphyxia in full term infants. *Arch Dis Child Fetal Neonatal Ed.* 1995; 72:F34-F38.
- (26) Toet MC, Hellström-Westas L, Groenendaal F, Eken P, de Vries LS . Amplitude integrated EEG 3 and 6 hours after birth in full term neonates with hypoxic-ischaemic encephalopathy. *Arch Dis Child Fetal Neonatal Ed.* 1999; 81:F19-F23.
- (27) Al Naqeeb N, Edwards AD, Cowan F, Azzopardi D. Assessment of neonatal encephalopathy by amplitude integrated electroencephalography. *Pediatrics* 1999; 103:1263-1271
- (28) Hellström-Westas L. Comparison between tape-recorded and amplitude integrated EEG monitoring sick newborn infants. *Acta Paediatr Scand* 1992; 81:812-819
- (29) Shalak LF, Laptook AR, Velaphi SC, Perlman JM. Amplitude-integrated electroencephalography coupled with an early neurologic examination enhances prediction of term infants at risk for persistent encephalopathy. *Pediatrics* 2003; 111:351-351.

- (30) Osredkar D, Toet MC, van Rooij LGM, van Huffelen AC, Groenendaal F, de Vries LS. Sleep-wake cycling on amplitude-integrated EEG in full-term newborns with hypoxic-ischemic encephalopathy. In press; *Pediatrics* Pediatrics, 2005;115:327-332
- (31) Van Rooij,L.G, Toet MC, Osredkar D, van Huffelen AC, Groenendaal F, de Vries LS. Recovery of amplitude integrated electroencephalographic background patterns within 24 hours of perinatal asphyxia. *Arch Dis Child Fetal Neonatal Ed.* 2005 May;90(3):F245-51
- (32) Sher M.S, Alvin J, Gaus L, Minnigh B, Painter M. J. Uncoupling of EEG-clinical neonatal seizures after antiepileptic drug use. *Pediatr Neurol* 2003; 28:277-280.
- (33) Boylan GB, Rennie JM, Pressler RM, Wilson G, Morton M, Binnie CD. Phenobarbitone, neonatal seizures, and video-EEG. *Arch Dis Child Fetal Neonatal Ed* 2002; 86: F165-F170.
- (34) Meierkord H, Boon, P, Engelsen B, Gocke K, Shorvon S, Tinuper P, Holtkamp M. EFNS guideline on the management of sttus epilepticus. *Eur J Neurol* 2006, May; 13(5):445-50.
- (35) Peets A, Berthiaume L, Bagshaw S, Frederico P, Doig Ch, Zygun D, Prolonged refractory status epilepticus following acute traumatic brain injury: a case report of excellent neurological recovery. *Critical Care* 2005, 9:R725-R728 (DOI 10.1186/cc3884)
- (36) Rosetti A, Reichhart M, Schaller M, Despland P, Bogousslavsky J, Propofol treatment of refractory status epilepticus: a study of 31 episodes, *Epilepsia*, 45(7): 757-763, 2004
- (37) Claassen J, Bäumer T, Hansen H, Kontinuierliches EEG zum Monitoring auf der Intensivstation: Neue Einsatzmöglichkeiten und Nutzen für therapeutische Entscheidungen, *Der Nervenarzt* Heft 71 Nr. 10, Sept. 2000